

## Acute intracerebroventricular injection of chemerin-9 increases systemic blood pressure via CMKLR1 in rats

Yamamoto Atsunori<sup>1</sup>, Kengo Matsumoto<sup>1</sup>, Satoshi Kameshima<sup>2</sup>, Naoko Yamaguchi<sup>3</sup>, Syousirou Okada<sup>3</sup>, Muneyoshi Okada<sup>1</sup>, Hideyuki Yamawaki<sup>1</sup>

<sup>1</sup>Lab. Vet Pharmacol., Sch., Vet. Med., Kitasato Univ., <sup>2</sup>Lab. Small Animal Internal Med., Sch., Vet. Med., Kitasato Univ., <sup>3</sup>Dept. Pharmacol., Sch Med., Aichi Med. Univ.

Chemerin is an inflammatory adipocytokine through acting on G protein-coupled receptor, chemokine-like receptor (CMKLR)1. We previously demonstrated that long-term intraperitoneal administration with chemerin increased systolic blood pressure (BP) in mice. Since important nuclei regulating BP exist in the brain, we examined mechanisms of central BP control by chemerin-9.

BP was invasively measured after acute intracerebroventricular (i.c.v.) injection of chemerin-9 to CMKLR1 siRNA-treated rats for 3 days. Serum adrenaline was measured by HPLC method. CMKLR1 expression in the brain tissues of spontaneously hypertensive rat (SHR) was examined by Western blotting.

In the control siRNA-treated rats, chemerin-9 significantly increased mean BP, while it had no effect in the CMKLR1 siRNA-treated rats. Serum adrenaline level was increased by the acute i.c.v. injection of chemerin-9. CMKLR1 expression around brain ventricles from SHR was increased.

We for the first time demonstrated that chemerin-9 stimulates the sympathetic nerves via CMKLR1 expressed around brain ventricles, which leads to an increase in BP. It is further suggested that chemerin/CMKLR1 in the central nerves plays an important role in essential hypertension.