

Loss of P2Y₁ receptor causes ocular hypertension and glaucoma-like phenotype in mice

Hamada Kentaro¹, Youichi Shinozaki¹, Takahiro Segawa², Kazuhiko Namekata³, Nobuhiko Ohno^{4,5}, Takayuki Harada³, Kenji Kashiwagi⁶, Schuichi Koizumi¹

¹Dept. Neuropharmacol., Interdiscip. Grad. Sch. Med. Univ. Yamanashi, ²Cent. Life Sci. Res., Univ. Yamanashi, ³Vis. Res. Project, Tokyo Metr. Inst. Med. Sci., ⁴Dev. Ultrastruct. Res., Natl. Inst. Physiol. Sci., ⁵Div. Anatomy, Jichi Med. Univ., ⁶Dept. Ophthalmol., Interdiscip. Grad. Sch. Med. Univ. Yamanashi

Glaucoma is second leading cause of blindness worldwide which is characterized by progressive degeneration of retinal ganglion cells (RGCs). Elevated intraocular pressure (IOP) is one of the highest risk factors and IOP-lowering agents are used to prevent glaucoma. New molecular target is required because of the side effects, drug resistance, and insufficiency for IOP reduction by a part of pre-existing agents. Here, we report that P2Y₁ receptor (P2Y₁R) activation induces IOP reduction and knock out of P2Y₁R (P2Y₁KO) causes sustained IOP elevation associated with age-dependent RGC degeneration. Topical application of MRS2365, selective agonist for P2Y₁R, caused significant reduction in IOP in wild-type (WT) mice but not in P2Y₁KO mice. We also found that P2Y₁KO mice showed significantly higher IOP level than that in WT mice. Because sustained IOP elevation is one feature of hypertensive glaucoma, we checked RGC damages and found that the number of RGCs in P2Y₁KO mice was comparable at 3 months old but significantly smaller at 12 months old. Furthermore, optical coherence tomography (OCT) revealed that 12-month-old P2Y₁KO mice showed thinner ganglion cell and inner plexiform layers, general diagnostic feature of glaucoma patients. Taken together, our results demonstrated that (1) P2Y₁R activation reduces IOP; (2) loss-of-function of P2Y₁R causes sustained elevation in IOP and (3) hypertensive glaucoma-like phenotypes in middle-aged mice.